

A NOTE ON THE SUMS OF POWERS OF CONSECUTIVE q -INTEGERS

YILMAZ SIMSEK, DAYEOUL KIM, TAEKYUN KIM, AND SEOG-HOON RIM

ABSTRACT. In this paper we construct the q -analogue of Barnes's Bernoulli numbers and polynomials of degree 2, which is an answer to a part of Schlosser's question. Finally, we will treat the q -analogue of the sums of powers of consecutive integers.

1. INTRODUCTION

In 1713, J. Bernoulli first discovered the method which one can produce those formulae for the sum $\sum_{j=1}^n j^k$, for any natural numbers k (cf. [1],[3],[6],[7],[15],[22]). The Bernoulli numbers are among the most interesting and important number sequences in mathematics. These numbers first appeared in the posthumous work “*Ars Conjectandi*” (1713) by Jakob Bernoulli(1654-1705) in connection with sums of powers of consecutive integers (Bernoulli(1713) or D. E. Smith(1959) see [15]).

Let q be an indeterminate which can be considered in complex number field, and for any integer k define the q -integer as

$$[k]_q = \frac{q^k - 1}{q - 1}, \text{ (cf. [11],[12],[13],[16],[17]).}$$

Note that $\lim_{q \rightarrow 1} [k]_q = k$. Recently, many authors studied q -analogue of the sums of powers of consecutive integers.

In [6], Garrett and Hummel gave a combinatorial proof of a q -analogue of $\sum_{k=1}^n k^3 = \binom{n+1}{2}^2$ as follows:

$$\sum_{k=1}^n q^{k-1} \left(\frac{1 - q^k}{1 - q} \right)^2 \left(\frac{1 - q^{k-1}}{1 - q^2} + \frac{1 - q^{k+1}}{1 - q^2} \right) = \left[\begin{matrix} n+1 \\ 2 \end{matrix} \right]_q^2,$$

where

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_q = \prod_{j=1}^k \frac{1 - q^{n+1-j}}{1 - q^j}$$

2000 AMS Subject Classification: 11B68, 11S40.

keywords and phrases :Sums of powers, Bernoulli Numbers, q -Bernoulli Numbers, zeta function, Dirichlet series

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

denotes the q -binomial coefficient. Garrett and Hummel, in their paper, asked for a simpler q -analogue of the sum of cubes. As a response to Garrett and Hummel's question, Warnaar gave a simple q -analogue of the sum of cubes as follows:

$$\sum_{k=1}^n q^{2n-2k} \frac{(1-q^k)^2 (1-q^{2k})}{(1-q)^2 (1-q^2)} = \left[\begin{matrix} n+1 \\ 2 \end{matrix} \right]_q^2. \quad (1.1)$$

In [21], Schlosser took up on Garrett and Hummel's second question. Especially, he studied the q -analogues of the sums of consecutive integers, squares, cubes, quarts and quints. He obtained his results by employing specific identities for very-well-poised basic hypergeometric series. However, Schlosser could not find the q -analogue of the sums of powers of consecutive integers of higher order, and left it as question. In [15], T. Kim evaluated sums of powers of consecutive q -integers as follows:

For any positive integers $n, k (> 1)$, $h \in \mathbb{Z}$, let

$$S_{n,q^h}(k) = \sum_{j=0}^n q^{hj} [k]_q^n.$$

Then, he obtained the interesting formula for $S_{n,q}(k)$ below:

$$S_{n,q}(k) = \frac{1}{n+1} \sum_{j=0}^n \binom{n+1}{j} \beta_{j,q} q^{kj} [k]_q^{n+1-j} - \frac{(1-q^{(n+1)k}) \beta_{n+1,q}}{n+1},$$

where $\beta_{j,q}$ are the modified Carlitz's q -Bernoulli numbers. Indeed, this formula is exactly a q -analogue of the sums of powers of consecutive integers due to Bernoulli.

Recently, the problem of q -analogues of the sums of powers have attracted the attention of several authors([8],[9],[15],[19],[21],[23]). Let

$$S_{m,n}(q) = \sum_{k=1}^n [k]_{q^2} [k]_q^{m-1} q^{(n-k) \frac{m+1}{2}}. \quad (1.2)$$

Then Warnaar[23] (for $m = 3$) and Schlosser[21] gave formulae for $m = 1, 2, 3, 4, 5$ as the meaning of the q -analogues of the sums of consecutive integers, squares, cubes, quarts and quints. By two families of polynomials and Vandermonde determinant, Guo and Zeng[9] found the formulae for the q -analogues of the sums of consecutive integers (for $m = 1, 2, \dots, 5$). They recovered the formulae of Warnaar and Schlosser, for $m = 6, 7, \dots, 11$. In [21], Schlosser speculated on the existence of a general formula for $S_{m,n}(q)$, which is defined in (1.2), and left it as an open problem.

By using T. Kim technical method to construct q -Bernoulli numbers and polynomials in [10],[11],[12],[13],[14],[15],[16],[19], we construct the q -analogue of Barnes' Bernoulli numbers and polynomials of degree 2, which is an answer to a part of Schlosser's question. Finally, we give some formulae for $S_{m,n}(q)$.

We define generating function $F_{k,q}^*(t)$ of the q -Bernoulli numbers $\beta_{n,k,q}^*$ ($n \geq 0$) as follows:

$$\begin{aligned} F_{k,q}^*(t) &= -t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \exp(t[j]_q q^{\frac{k-j}{2}}) \\ &= \sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!} \quad (\text{cf. [10], [11], [12], [13], [14], [15], [16]}). \end{aligned} \quad (1.3)$$

Theorem 1. *Let n, k be positive integers. Then*

$$\beta_{n,k,q}^* = \left(\frac{1}{1-q} \right)^{n-1} \sum_{m=0}^n \binom{n}{m} \frac{(-1)^m m q^{\frac{(n-1)(k-1)}{2} + k + m - 2}}{(1 - q^{m - \frac{n-1}{2} - 2})(1 - q^{m - \frac{n-1}{2}})}.$$

We define generating function $F_{k,q}^*(t; k)$ of the q -Bernoulli polynomials $\beta_{n,k,q}^*(k)$ ($n \geq 0$) as follows:

$$F_{k,q}^*(t; k) = -t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} \exp(t[j+k]_q q^{\frac{-j}{2}}) = \sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k) t^n}{n!}.$$

Theorem 2. *Let n, k be positive integers. Then*

$$\beta_{n,k,q}^*(k) = \frac{1}{[2]_q (1-q)^{n-2}} \sum_{m=0}^n \binom{n}{m} (-1)^m \left(\frac{m q^{k(m-1)}}{1 - q^{m - \frac{n-1}{2} - 2}} - \frac{m q^{k(m+1)}}{1 - q^{m - \frac{n-1}{2}}} \right).$$

Theorem 3. (General formula for $S_{m,n}(q)$) *Let n, k be positive integers. Then*

$$S_{n,k}(q) = \sum_{j=0}^{k-1} [j]_{q^2} [j]_q^{n-1} q^{\frac{(n+1)(k-j)}{2}} = \frac{\beta_{n,k,q}^*(k) - \beta_{n,k,q}^*}{n}.$$

2. PRELIMINARY

Let n, k be positive integers ($k > 1$), and let

$$S_n(k) = \sum_{j=1}^{k-1} j^n = 1^n + 2^n + \dots + (k-1)^n.$$

It is well-known that

$$\begin{aligned} S_1(k) &= \frac{1}{2}k^2 - \frac{1}{2}k, \\ S_2(k) &= \frac{1}{3}k^3 - \frac{1}{2}k^2 + \frac{1}{6}k, \\ S_3(k) &= \frac{1}{4}k^4 - \frac{1}{2}k^3 + \frac{1}{4}k^2, \\ &\dots \end{aligned}$$

Thus we have the following three conjectures:

(I) $S_n(k)$ is a polynomial in k of degree $n + 1$ with leading coefficient $\frac{1}{n+1}$,

(II) The constant term of $S_n(k)$ is zero, i.e., $S_n(0) = 0$,

(III) The coefficient of k^n in $S_n(k)$ is $-\frac{1}{2}$.

Therefore, $S_n(k)$ is a polynomial in k of the form

$$S_n(k) = \frac{1}{n+1}k^{n+1} - \frac{1}{2}k^n + a_{n-1}k^{n-1} + \cdots + a_1k.$$

We note that

$$\frac{d}{dk}S_n(k) = k^n - \frac{n}{2}k^{n-1} + \cdots.$$

To make life easier, we put the first two conjectures together and we reach the following conjecture, which is what Jacques Bernoulli (1654-1705) claimed more than three hundred years ago.

Bernoulli: There exists a unique monic polynomial of degree n , say $B_n(x)$, such that

$$S_n(k) = \sum_{j=1}^{k-1} j^n = 1^n + 2^n + \cdots + (k-1)^n = \int_0^k B_n(x)dx.$$

As the q -analogue of $S_n(k)$, Schlosser[21] considered the existence of general formula on $S_{m,n}(q)$, and he gave the below values:

$$\sum_{k=1}^n [k]_{q^2} [k]_q^{m-1} q^{(n-k)\frac{m+1}{2}},$$

where $m = 1, 2, \dots, 5$.

Indeed,

$$\sum_{k=1}^n [k]_{q^2} [k]_q q^{\frac{3}{2}(n-k)} = \frac{[n]_q [n+1]_q [n+\frac{1}{2}]_q}{[1]_q [2]_q [\frac{3}{2}]_q}, \quad m = 2,$$

$$\sum_{k=1}^n [k]_{q^2} [k]_q^2 q^{2(n-k)} = \left[\begin{matrix} n+1 \\ 2 \end{matrix} \right]_q^2, \quad m = 3,$$

$$\begin{aligned} \sum_{k=1}^n [k]_{q^2} [k]_q^3 q^{\frac{5}{2}(n-k)} &= \frac{(1-q^n)(1-q^{n+1})(1-q^{n+\frac{1}{2}})}{(1-q)(1-q^2)(1-q^{\frac{5}{2}})} \\ &\times \left(\frac{(1-q^n)(1-q^{n+1})}{(1-q)^2} - \frac{q^n(1-q^{\frac{1}{2}})}{1-q^{\frac{3}{2}}} \right), \quad m = 4, \end{aligned}$$

$$\begin{aligned} \sum_{k=1}^n [k]_{q^2} [k]_q^4 q^{3(n-k)} &= \frac{(1-q^n)^2(1-q^{n+1})^2}{(1-q)^2(1-q^2)(1-q^3)} \\ &\times \left(\frac{(1-q^n)(1-q^{n+1})}{(1-q)^2} - \frac{q^n(1-q)}{1-q^2} \right), \quad m = 5. \end{aligned}$$

T. Kim[15] proved the smart formula for the q -analogue of $S_n(k)$ as follows:

$$\sum_{k=0}^{n-1} q^k [k]_q = \frac{1}{2} \left([n]_q^2 - \frac{[2n]_q}{[2]_q} \right)$$

and

$$\sum_{k=0}^{n-1} q^{k+1} [k]_q^2 = \frac{1}{3} [n]_q^3 - \frac{1}{2} \left([n]_q^2 - \frac{[2n]_q}{[2]_q} \right) - \frac{1}{3} \frac{[3n]_q}{[3]_q}.$$

3. PROOF OF MAIN THEOREMS

We define generating function $F_{k,q}^*(t)$ of the q -Bernoulli numbers $\beta_{n,k,q}^*$ ($n \geq 0$) as follows:

$$F_{k,q}^*(t) = -t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \exp(t[j]_q q^{\frac{k-j}{2}}) = \sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!}. \quad (3.1)$$

Proof of Theorem 1. Let

$$\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!} = -t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \exp(t[j]_q q^{\frac{k-j}{2}}).$$

By using Taylor series in the above then we have

$$\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!} = -t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \sum_{n=0}^{\infty} \frac{[j]_q^n q^{\frac{n(k-j)}{2}}}{n!} t^n.$$

By using some elementary calculations in the above, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!} &= -t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \\ &\quad \times \sum_{n=0}^{\infty} \left\{ \left(\frac{1}{1-q} \right)^n q^{\frac{n(k-j)}{2}} \sum_{m=0}^n \binom{n}{m} (-1)^m q^{jm} \right\} \frac{t^n}{n!} \\ &= \frac{-t}{q^2 - 1} \sum_{n=0}^{\infty} \left(\frac{1}{1-q} \right)^n q^{k + \frac{nk}{2}} \sum_{m=0}^n \binom{n}{m} (-1)^m \\ &\quad \times \sum_{j=0}^{\infty} \left(q^{mj-j-\frac{jn}{2}} (1 - q^{2j}) \right) \frac{t^n}{n!}. \end{aligned}$$

By using geometric power series in the above and after some calculations, we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!} &= \frac{-t}{q^2 - 1} \sum_{n=0}^{\infty} \left(\frac{1}{1-q} \right)^n \\ &\quad \times \sum_{m=0}^n \binom{n}{m} \frac{(-1)^m q^{m+k+\frac{n}{2}(k-1)-1} (1-q^2)}{(1-q^{-1+m-\frac{n}{2}}) (1-q^{1+m-\frac{n}{2}})} \frac{t^n}{n!}. \end{aligned}$$

For $n = 0$, then $\beta_{0,k,q}^* = 0$. Thus, we have

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\beta_{n,k,q}^* t^n}{n!} &= -t \sum_{n=1}^{\infty} \left(\frac{1}{1-q} \right)^{n-1} \\ &\quad \times \sum_{m=1}^{n-1} \binom{n-1}{m-1} \frac{(-1)^{m-1} q^{m+k+\frac{(n-1)(k-1)}{2}-2}}{(1-q^{-2+m-\frac{n-1}{2}}) (1-q^{m-\frac{n-1}{2}})} \frac{t^{n-1}}{(n-1)!} \\ &= \sum_{n=1}^{\infty} \left(\frac{1}{1-q} \right)^{n-1} \\ &\quad \times \sum_{m=0}^n \binom{n}{m} \frac{(-1)^m m q^{m+k+\frac{(n-1)(k-1)}{2}-2}}{(1-q^{m-\frac{n-1}{2}-2}) (1-q^{m-\frac{n-1}{2}})} \frac{t^n}{n!}. \end{aligned}$$

By comparing the coefficients of $\frac{t^n}{n!}$ on both sides of the above equation, we easily arrive at the desired result. $\square$

We define generating function $F_{k,q}^*(t; k)$ of the q -Bernoulli polynomials $\beta_{n,k,q}^*(k)$ ($n \geq 0$) as follows:

$$F_{k,q}^*(t; k) = -t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} \exp(t[j+k]_q q^{\frac{-j}{2}}) = \sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k) t^n}{n!}. \quad (3.2)$$

Proof of Theorem 2. Let

$$\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k) t^n}{n!} = -t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} \exp(t[j+k]_q q^{\frac{-j}{2}}).$$

By using Taylor expansion of e^x in the above, we have

$$\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k) t^n}{n!} = -t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} \sum_{n=0}^{\infty} \frac{[j+k]_q^n q^{\frac{-nj}{2}}}{n!} t^n.$$

By using some elementary calculations in the above, we have

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k)t^n}{n!} &= -t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} \\
&\times \sum_{n=0}^{\infty} \left\{ \left(\frac{1}{1-q} \right)^n \sum_{m=0}^n \binom{n}{m} (-1)^m q^{m(j+k) - \frac{nj}{2}} \right\} \frac{t^n}{n!} \\
&= \frac{t}{q^2-1} \sum_{n=0}^{\infty} \left(\frac{1}{1-q} \right)^n \sum_{m=0}^n \binom{n}{m} (-1)^m q^{mk} \\
&\times \sum_{j=0}^{\infty} \left(1 - q^{2(j+k)} \right) q^{-j+jm - \frac{jn}{2}} \frac{t^n}{n!}.
\end{aligned}$$

By using geometric power series in the above and after some calculations, we obtain

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k)t^n}{n!} &= -t \sum_{n=0}^{\infty} \frac{1}{[2]_q} \left(\frac{1}{1-q} \right)^{n-1} \\
&\times \sum_{m=0}^{n-1} \binom{n}{m} \frac{(-1)^{m+1} q^{\frac{n-1}{2}(k-1)+k+m-1}}{\left(1 - q^{-1+m-\frac{n-1}{2}} \right) \left(1 - q^{1+m-\frac{n-1}{2}} \right)} \frac{t^{n-1}}{(n-1)!}.
\end{aligned}$$

Thus, we get

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{\beta_{n,k,q}^*(k)t^n}{n!} &= -t \sum_{n=0}^{\infty} \frac{1}{[2]_q} \left(\frac{1}{1-q} \right)^{n-1} \\
&\times \sum_{m=0}^n \binom{n}{m} (-1)^m \left(\frac{q^{mk}}{1 - q^{m-1-\frac{n}{2}}} - \frac{q^{(m+2)k}}{1 - q^{m+1-\frac{n}{2}}} \right) \frac{t^n}{n!}
\end{aligned}$$

and

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{\beta_{n,k,q}^*(k)t^n}{n!} &= \sum_{n=1}^{\infty} \frac{1}{[2]_q} \left(\frac{1}{1-q} \right)^{n-2} \\
&\times \sum_{m=0}^n \binom{n}{m} (-1)^m \left(\frac{mq^{(m-1)k}}{1 - q^{m-\frac{n-1}{2}-2}} - \frac{mq^{(m+1)k}}{1 - q^{m-\frac{n-1}{2}}} \right) \frac{t^n}{n!}.
\end{aligned}$$

By comparing the coefficients of $\frac{t^n}{n!}$ on both sides of the above equations, we easily arrive at the desired result. $\square$

Proof of Theorem 3. Let

$$\begin{aligned}
& - \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} \exp(t[j+k]_q q^{\frac{-j}{2}}) + \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \exp(t[j]_q q^{\frac{k-j}{2}}) \\
&= \sum_{j=0}^{k-1} q^{k-j} [j]_{q^2} \exp(t[j]_q q^{\frac{k-j}{2}}).
\end{aligned} \tag{3.3}$$

By using Theorem 1 and Theorem 2 and (3.3), we easily arrive at the desired result. Our proof of Theorem 3 runs parallel that of Theorem 1 and Theorem 2 above, so we choose to omit the details involved.

Remark 1. The Barnes double zeta function is defined by

$$\zeta_2(s, w \mid w_1, w_2) = \sum_{m, n=0}^{\infty} (w + mw_1 + nw_2)^{-s}, \quad \text{Re}(s) > 2, \quad (\text{cf. [2]}),$$

for complex number $w \neq 0, w_1, w_2$ with positive real parts.

The Barnes' polynomial: Barnes[2] introduced r -tuple Bernoulli polynomials ${}_rS_m(u; \tilde{w})$ by,

$$\begin{aligned} F_r(t; u; \tilde{w}) &= \frac{(-1)^r t e^{-ut}}{\prod_{j=1}^r (1 - e^{-w_j t})} = \sum_{k=1}^r \frac{{}_rS_1^{(k+1)}(u; \tilde{w}) (-1)^k}{t^{k-1}} \\ &+ \sum_{m=1}^{\infty} \frac{{}_rS'_m(u; \tilde{w})}{m!} (-1)^{m-1} t^m, \end{aligned} \quad (3.4)$$

for $|t| < \min \left\{ \frac{2\pi}{|w_1|}, \dots, \frac{2\pi}{|w_r|} \right\}$. Here $w_1, w_2, \dots, w_r$ are complex number with positive real parts, $\tilde{w} = (w_1, w_2, \dots, w_r)$ and ${}_rS_1^{(k)}(u; \tilde{w})$ the k -th derivative of ${}_rS_1(u; \tilde{w})$ with respect to u .

Substituting $u = -1$, $w_1 = w_2 = -1$ and $r = 2$ into (3.4), we have

$$F_2(t; -1; -1, -1) = \frac{te^t}{(1 - e^t)^2}.$$

In (3.1),

$$\lim_{q \rightarrow 1} F_{k,q}^*(t) = \lim_{q \rightarrow 1} -t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} \exp(t[j]_q q^{\frac{k-j}{2}}) = -F_2(t; -1; -1, -1).$$

4. FURTHER REMARKS AND OBSERVATIONS ON A CLASS OF q - ZETA FUNCTIONS

In this section, by using generating functions of $F_{k,q}^*(t)$ and $F_{k,q}^*(t; k)$, we produce new definitions of q -polynomials and numbers. These generating functions are very important in the case of defining q -zeta function. Therefore, by using these generating functions and Mellin transformation, we will define the q -zeta function.

By applying Mellin transformation in (3.2), we obtain

$$\frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2} F_{k,q}^*(-t; k) dt = \sum_{n=0}^{\infty} \frac{[n+k]_{q^2} q^{-n \frac{s+2}{2}}}{[n+k]_q^s},$$

where $\Gamma(s)$ denotes the Euler gamma function.

For $s \in \mathbb{C}$, we define

$$\zeta_{k,q}^*(s; k) = \sum_{n=0}^{\infty} \frac{[n+k]_q^2 q^{-n \frac{s+2}{2}}}{[n+k]_q^s}, \quad \operatorname{Re}(s) > 2.$$

By Mellin transformation in (3.1), we obtain

$$\begin{aligned} \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2} F_{k,q}^*(-t) dt &= \sum_{n=0}^{\infty} \frac{[n]_q^2 q^{\frac{(k-n)(2-s)}{2}}}{[n]_q^s} \\ &= \zeta_{k,q}^*(s), \operatorname{Re}(s) > 2. \end{aligned}$$

For any positive integer n , Cauchy Residue Theorem in the above equation, we have

$$\zeta_{k,q}^*(1-n) = -\frac{\beta_{n,k,q}^*}{n}.$$

REFERENCES

- [1] T. Apostol, *Introduction to analytic number theory*, Springer -Verlag, New York, 1976.
- [2] E. W. Barnes, *On theory of the multiple gamma functions*, Trans. Camb. Philos. Soc. **19** (1904), 374-425.
- [3] L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
- [4] E. Deeba and D. Rodriguez, *Stirling's series and Bernoulli numbers*, Amer. Math. Monthly **98** (1991), 423-426.
- [5] J. Faulhaber, *Academia Algebrae, Darinnen die miraculosische Inventiones zu den höchsten Cossen weiters continuirt und profitiert werden*, Augspurg, bey Johann Ulrich Schönnigs, 1631.
- [6] K. C. Garrett and K. Hummel, *A combinatorial proof of the sum of q-cubes*, Electron. J. Combin. **11** (2004).
- [7] G. Gasper and M. Rahman, *Basic hypergeometric series, Encyclopedia of Mathematics and Its Applications 35*, Cambridge University Press, Cambridge, 1990.
- [8] I. P. Goulden and D. M. Jackson, *Combinatorial enumeration, reprint of the 1983 original*, Dover Publications, Inc., Mineola, NY, 2004.
- [9] V. J. W. Guo and J. Zang, *A q-analogue of Faulhaber's formula for sums of powers*, arXiv:math.CO/0501441 (2005).
- [10] T. Kim, *An invariant p-adic integral associated with Daehee Numbers*, Integral Transforms Spec. Funct. **13** (2002), 65-69.
- [11] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **19** (2002), 288-299.
- [12] T. Kim, *On p-adic q-L-functions and sums of powers*, Discrete Math. **252** (2002), 179-187.
- [13] T. Kim, *Non-archimedean q-integrals associated with multiple Changhee q-Bernoulli Polynomials*, Russ. J. Math. Phys. **10** (2003), 91-98.
- [14] T. Kim, *On Euler-Barnes multiple zeta functions*, Russ. J. Math. Phys. **10** (2003), 261-267.
- [15] T. Kim, *Sums of powers of consecutive q-integers*, Adv. Stud. Contemp. Math. **9** (2004), 15-18.
- [16] T. Kim, *p-adic q-integrals associated with the Changhee-Barnes' q-Bernoulli polynomials*, Integral Transforms Spec. Funct. **15** (2004), 415-420.
- [17] T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russ. J. Math. Phys. **11** (2004), 71-76.
- [18] T. Kim, *A new approach to q-zeta function*, arXiv:math.NT/0502005 (2005).
- [19] T. Kim, *q-analogue of the sums of powers consecutive integers*, arXiv:math.NT/0502113 (2005).

- [20] D. E. Knuth, Johann Faulhaber and sums of powers, *Math. Comp.* **61** (1993), 277-294.
- [21] M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts and quints*, *Electron. J. Combin.* **11** (2004), #R71.
- [22] Y. -Y. Shen, *A note on the sums of powers of consecutive integers*, *Tunghai Science* **5** (2003), 101-106.
- [23] S. O. Warnaar, *On the q-analogue of the sum of cubes*, *Electron. J. Combin.* **11** (2004), #N13.
- [24] L. C. Washington, *Introduction to cyclotomic fields*, Springer-Verlag, New York, 1997.
- [25] E. T. Whittaker and G. N. Watson, *A course of modern analysis*, Cambridge University Press, London and New York, 1927.

YILMAZ SIMSEK

Mersin University, Faculty of Science,
Department of Mathematics 33343 Mersin, Turkey
e-mail: ysimsek@mersin.edu.tr

Daeyeoul Kim

Department of Mathematics,
Chonbuk National University,
Chonju, 561-756, Korea
e-mail: dykim@math.chonbuk.ac.kr

TAEKYUN KIM

Institute of Science Education,
Kongju National University Kongju 314-701, Korea
e-mail: tkim@kongju.ac.kr

Seog-Hoon Rim

Department of Mathematics Education,
Kyungpook National University, Taegu 702-701, Korea
e-mail : shrim@knu.ac.kr